

IIII Integrated Rate Laws

- From initial concentrations & rate law, we can predict all concentrations at any time *t*.
- Mathematically, this is an initial value problem involving a (usually) simple differential equation.





$^{3}_{1}H$	12.3 y	²³⁵ ₉₂ U	7.1×10^{8} y
¹⁴ ₆ C	$5.73 \times 10^{3} \text{ y}$	²³⁸ ₉₂ U	4.5×10 ⁹ y
¹⁵ ₆ C	2.4 s	¹³⁷ ₅₅ Cs	30.17
$^{40}_{19}$ K	1.26×10 ⁹ y	$^{131}_{53}$ I	8.05 d
⁹⁰ ₃₈ Sr	28.1 y	²²⁶ ₈₈ Ra	1.60×10^{3}
⁶⁰ ₂₇ Co	5.26 y		

🛄 Example

Hydrogen peroxide decomposes into water and oxygen in a first-order process.

$$\mathrm{H}_{2}\mathrm{O}_{2}(aq) \rightarrow \mathrm{H}_{2}\mathrm{O}(l) + \frac{1}{2}\mathrm{O}_{2}(g)$$

At 20.0 °C, the ½-life for the reaction is 3.92×10^4 seconds. If the initial concentration of hydrogen peroxide is 0.52 M, what is the concentration after 7.00 days (6.048 × 10⁵ s)?































